

Exotic neutron-rich nuclei in a three-body model

L. Tomio¹, T. Frederico², and M. T. Yamashita³

¹ Instituto de Física Teórica, Universidade Estadual Paulista
01405-900 São Paulo, Brazil

² Departamento de Física, Instituto Tecnológico de Aeronáutica, Centro Técnico
Aeroespacial 12228-900 São José dos Campos, Brazil

³ Unidade Diferenciada de Itapeva, Universidade Estadual Paulista
18409-010 Itapeva, Brazil

Received 1 January 2004

Abstract. We present results for spatial distributions of weakly-bound three-body systems, derived from a universal scaling function that depends on the mass ratio of the particles, as well as on the nature of the subsystems.

Keywords: few-body, root-mean-square radii, halo nuclei, correlation

PACS: 21.45.+v, 27.20.+n, 25.75.Gz

The approach to obtain the mean-square distances between particles in a weakly-bound three-body system [1,2] considers a universal scaling function that depends on the mass ratio of the neutron and the core, as well as on the nature of the subsystems, bound or virtual. In the model, we have a minimal number of physical inputs, directly related to observables. As considered in Ref. [1], for halo-nuclei systems consisting of a core with two halo neutrons, for the low-energy observables it was used the neutron-neutron and neutron-core s-wave scattering lengths (or the corresponding virtual or bound energies).

A generic weakly three-body system $\alpha\alpha\beta$ can be described according to the two-body subsystems properties, as discussed in Ref. [1]. We have four possible configurations for such a system: *Borromean* configuration, when all the two-body subsystems are unbound; *Tango* configuration, when we have two unbound and one bound two-body subsystems; *Samba* configuration, when just one two-body subsystem is unbound; and *All-bound* configuration, when there is no unbound subsystems. To examine such weakly bound systems, the corresponding Faddeev formalism is given in some detail in another contribution to this proceedings [3], where is also given some results for the size of a few halo-nuclei systems.

In the present contribution we resume in Fig.1 our results for the mean-square distances of generic weakly-bound three-body systems $\alpha\alpha\beta$ where two particles are

identical. In Figure 1, we have the root-mean-square (rms) distances between the particles in units of the inverse of the three-body binding energy E_3 , plotted as functions of $K_{\alpha\beta}/|K_{\alpha\alpha}|$ and $K_{\alpha\alpha}/|K_{\alpha\beta}|$, where $K_{\alpha\gamma} \equiv \pm\sqrt{E_{\alpha\gamma}/E_3}$ ($\gamma = \alpha, \beta$) and $E_{\alpha\gamma}$ is the energy of the $\alpha\gamma$ subsystem, with $+$ standing for bound and $-$ for unbound (virtual). Our units are such that $\hbar = 1$ and $m_\alpha = 1$. As evidenced by the results, where it was fixed E_3 , the size of the three-body systems are such that the smallest one is the one where all the two-body subsystems are unbound (Borromean).




Fig. 1. Dimensionless root-mean-square (rms) distances between the particles $\alpha\alpha$ and $\alpha\beta$, in units of the inverse of the three-body binding energy E_3 , given as functions of $K_{\alpha\gamma} \equiv \pm\sqrt{E_{\alpha\gamma}/E_3}$ ($\gamma = \alpha, \beta$), as explained in the text. In all the plots the mass ratio is such that $m_\beta/m_\alpha = 200$. In the left hand side we have $E_{\alpha\alpha}/E_3 = 0.1$; and, in the right hand side, $E_{\alpha\beta}/E_3 = 0.1$. The behavior or the results follows the same trend for other mass ratios and two-body energy parameters.

We thank partial support received from FAPESP and CNPq. MTY thanks support from the organizers of SLAFNAP6.

References

1. M.T. Yamashita, L. Tomio and T. Frederico, *Nucl. Phys. A* **735** (2004) 40.
2. M.T. Yamashita, T. Frederico and Lauro Tomio, *Phys. Rev. C* **72** (2005) 011601.
3. M. T. Yamashita, T. Frederico, and L. Tomio, *Spatial characteristics of borromean, tango, samba and all-bound halo nuclei*, contribution to this Proceedings.